

ACUTE ABDOMINAL PAIN IN CHILDREN DIFFERENTIAL DIAGNOSES

SYMPTOMS:	CONSIDER:
Severe colicky bouts of pain in child aged < 2 years, child may appear pale, quiet and unwell between bouts of pain +/- redcurrant stool	 Pyloric Stenosis (3-12 weeks M>F) Intussusception Volvulus
Baby +/- 6/52 age with projectile vomiting	Pyloric stenosis
Green, bile stained vomiting	ObstructionIntussusceptionileus
Diarrhoea with blood	 bacterial dysentery irritable bowel disease (IBD) Henloch-Schoenlein purpura (HSP) (NB abdo pain may precede rash) Haemolytic Uraemic Syndrome (HUS) gastroenteritis
Incomplete history, possible signs of bruising	Consider NAI
High fever, Loin pain, Dysuria Urine positive for nitrites and leucocytes	• UTI
Severe abdominal pain Vomiting Fever Tachycardia Subjective tenderness	 Appendicitis (NB typical history may be absent in younger children)
Swollen /inflamed/painful testis	Torsion of testesTrauma
Urine positive for ketones and glucose	DKA
Urine positive for blood	 Kidney stones Renal trauma HSP nephritis
Lowish Hb for age	• IBD
Raised platelets	trauma
Raised amylase	pancreatitis
Raised liver enzymes	hepatitisbiliary pathology
Teenage girl	 pregnancy ?ectopic Mittelschmerz Pelvic inflammatory disease
Afro-Caribbean background	sickle cell
Recent URTI	mesenteric adenitispneumonia